CONTRIBUTIONS REVISTA MEXICANA DE FÍSICA **49 SUPLEMENTO 2**, 145–147 OCTUBRE 2003

arXiv:quant-ph/0302189v2 7 Nov 2003

On susy periodicity defects of the Lamé Potentials

Oscar Rosas-Ortiz
Departamento de Física, CINVESTAV-IPN, A.P. 14-740, 07000 México D.F., Mexico

Recibido el 28 de marzo de 2002; aceptado el 19 de junio de 2002

The supersymmetric Quantum Mechanics approach is applied to embed bound states in the energy gaps of periodic potentials. The mechanism to generate periodicity defects in the first Lamé potential is analyzed. The related bound states are explicitly derived.

Let $H=-\frac{d^2}{dx^2}+V(x)$ be a Schrödinger operator with $V(x)$ a real-valued and locally bounded function on $\mathbb{R}$. The spectrum $\sigma(H)$ of H, as well as its resolvent set $\rho(H)$, will be then complementary subsets of $\mathbb{R}$. The involved eigenvalue equation $H\,\psi(x)=\epsilon\psi(x)$, admits solutions for arbitrary $\epsilon\in\mathbb{R}$. The ϵ-values such that $\epsilon=E\in\sigma_d(H)$ lead to square integrable wave-functions $\psi_E(x)$ in $L^2(\mathbb{R})$, while for $\epsilon\in\rho(H)$ one gets eigenfunctions $\psi(x)$ deprived of physical meaning. It is well known that the class of Schrödinger's operators H admitting analytic expressions for the corresponding eigenfunctions is narrow. The supersymmetric (susy) Quantum Mechanics [1] intends to expand the class. In particular, the conventional Infeld-Hull factorization method has been extended to employ not only the physically interpretable ψ_E but also the nonphysical eigenfunctions ψ [2]. The main idea of this generalized factorization is to implement the Darboux-Crum transformation as the mechanism to construct multi-parametric families $\{V(\lambda_1,\lambda_2,...,\lambda_n,x)\}$ of (almost) isospectral exactly solvable potentials [3, 4]. The members of each one of these families share common analytical properties, such as the number and type of (possible) singularities as well as the asymptotic behaviour. Hence, it is intended that the procedure does not add new singularities nor modifies the asymptotic behaviour of the initial potentials. This mechanism is ensured by taking eigenfunctions ψ with $\epsilon\in\rho(H)$ as the key transformation functions [4].

The present contribution deals with the generalized factorization method applied to periodic potentials. As we shall see, the very interesting phenomenon of periodicity defects in crystal lattices can be produced as a consequence of the susy mechanism. We support our analysis mainly on the results reported recently in [5].

Let us consider a Schrödinger operator H_0 for which the solutions $u(x,\epsilon)$ to the eigenvalue equation

$$H_0u(x,\epsilon)=\epsilon u(x,\epsilon) \tag{1}$$

are known for any $\epsilon\in\mathbb{R}$. Notice that equation (1) does not require the square integrability of $u(x,\epsilon)$. An important example is the Hamiltonian with a real-valued, continuous and periodic potential

$$V_0(x+\tau)=V_0(x),\qquad x\in\mathbb{R}. \tag{2}$$

In this case, the translation operator $Tu(x,\epsilon)=u(x+\tau,\epsilon)$, maps the two-dimensional subspace $\mathcal{H}_\epsilon$ of solutions of the eigenvalue equation (1) into itself. As a consequence, there is a basis of eigenfunctions $\{\mathrm{v}_1(x),\mathrm{v}_2(x)\}$ in $\mathcal{H}_\epsilon$ for which T admits a 2×2 matrix representation $[T]$ (the *monodromy matrix*). The eigenvalue equation for $[T]$, evaluated at $x=T$, determines two eigenvalues $t_\pm$ for a given energy ϵ, i.e., $t_\pm=t_\pm(\epsilon)$. The corresponding eigenfunctions $v_\pm(x,\epsilon)$ exist for any $\epsilon\in\mathbb{R}$ and are called *Bloch functions*. The structure of $v_\pm(x,\epsilon)$ depends essentially on the values of ϵ, namely, if ϵ is such that $t_\pm\in S^1$, then $\epsilon\in\sigma(H_0)$ and the related Bloch functions become complex and bounded. The spectrum of H_0 is then continuous, composed by a series of spectral bands in $\mathbb{R}$. The edges of that bands are given by the ϵ-values such that $t_\pm=\pm1$, and the Bloch functions become τ-periodic or τ-antiperiodic. Finally, if ϵ is such that $t_\pm\in\mathbb{R}/\{\pm1\}$, then $\epsilon\in\rho(H_0)$ and the related Bloch functions are real and diverge at either $-\infty$ or $+\infty$. Due to this mechanism, $\rho(H_0)$ splits into a (finite or infinite) sequence of forbidden gaps. In summary, $\sigma(H_0)=\{\epsilon\,;\,t_\pm\in S^1\}$, while $\rho(H_0)=\mathbb{R}/\sigma(H_0)$.

As regards to the supersymmetric treatment of periodic Hamiltonians, one can consider two different options: (i) The Darboux transformations are such that, when applied to systems described by (2), the new potentials are also τ-periodic. (ii) The procedure does not respect the global periodicity of the initial potential. In the former case, Dunne and Feinberg [6] have shown that, sometimes, the nonsingular first order transformed potential becomes simply a half-period displaced copy of the initial one. Such potentials were called *self-isospectral*. A further study (using the band edge eigenfunctions) produced the 2-susy self-isospectral potentials [7]. The above results have been recently generalized to embrace arbitrary δ-displaced copies of the initial periodic potential [5]. Such a phenomenon has been named the *translational Darboux invariance* (or simply Darboux invariance), and it reproduces the self-isospectral case for $\delta=\tau/2$. The Darboux invariance is achieved by using Bloch solutions as transformation functions for ϵ either at the band edges or in the resolvent set of the periodic Hamiltonian H_0. Concerning the option (ii), we first remark that arbitrary linear combinations of $v_\pm(x,\epsilon)$ for $\epsilon\in\rho(H_0)$, when used as transformation functions $u(x,\epsilon)$ can produce regular non-periodic modifications in the periodic potential (2). As we are going to show, moreover, the functions

$u(x,\epsilon)$ allow to embed discrete energy levels in the gaps of H_0 and to determine the corresponding square integrable eigenfunctions [5].

The *generalized* first order factorization method [2] allows to obtain the solutions $\phi_{\epsilon'}(x,\epsilon)$ of a transformed eigenvalue equation $H_1(\epsilon)\phi_{\epsilon'}(x,\epsilon) = \epsilon'\phi_{\epsilon'}(x,\epsilon)$; $H_1(\epsilon) := -\frac{d^2}{dx^2}+V_1(x,\epsilon)$, where the new potential $V_1(x,\epsilon)$ is related with the initial one $V_0(x)$ by means of a Darboux transformation

$$V_1(x,\epsilon) = V_0(x) + 2\beta'(x,\epsilon). \tag{3}$$

The function β here is nothing but the general solution of the Riccati equation

$$-\beta'(x,\epsilon) + \beta^2(x,\epsilon) = V_0(x) - \epsilon. \tag{4}$$

At this stage, it is interesting to notice that the transformation $\beta(x,\epsilon) \equiv -\frac{d}{dx}\ln u(x,\epsilon)$ allows to simplify the expressions for the potential (3) and the involved eigenfunctions $\phi_{\epsilon'}(x,\epsilon)$ leading to:

$$V_1(x,\epsilon) = V_0(x) - 2\frac{d^2}{dx^2}\ln u(x,\epsilon),$$

$$\phi_{\epsilon'}(x,\epsilon) \propto \frac{W(u,\psi_{\epsilon'})}{u(x,\epsilon)}$$

where, by simplicity, we have written $u \equiv u(x,\epsilon)$ and $\psi_{\epsilon'} \equiv \psi_{\epsilon'}(x)$. The symbol $W(u,\psi_{\epsilon'})$ represents the Wronskian of u and $\psi_{\epsilon'}$.

The analytical behaviour of $\phi_{\epsilon'}(x,\epsilon)$ strongly depends on the values of ϵ and ϵ'. If $\epsilon' = E \in \sigma(H_0)$ and $E_0 > \epsilon \in \rho(H_0)$, with $E_0 \in \sigma(H_0)$ denoting the lowest band edge, the eigenfunction $\phi_E(x,\epsilon)$ can be bounded. A particular detail of the method is that the function $\phi_\epsilon(x,\epsilon) \propto 1/u(x,\epsilon)$ satisfies identically $H_1(\epsilon)\phi_\epsilon(x,\epsilon) = \epsilon\phi_\epsilon(x,\epsilon)$ and becomes orthogonal to all the $\phi_E(x,\epsilon)$. Moreover, it can be chosen in $L^2(\mathbb{R})$. Thereby $\sigma(H_1(\epsilon)) = \sigma(H_0)\cup\{\epsilon\}$. In other words, the Hamiltonian $H_1(\epsilon)$ can be constructed in order to have a bound state embedded in the lowest spectral gap of H_0. Hence, $H_1(\epsilon)$ will be almost isospectral with H_0 for a properly selected ϵ. This situation allows to consider them as susy partner Hamiltonians. Concerning the analytical properties of $V_1(x,\epsilon)$, it is very common to find that those conditions on ϵ, making $\phi_E(x,\epsilon)$ a square integrable function, lead to potentials (3) with exactly the same number and nature of divergences as the initial one (see for example [4]). In particular, as for $\epsilon < E_0$ the function $u(x,\epsilon)$ can be chosen nodeless, the potential $V_1(x,\epsilon)$ in (5) is just a regular non-periodic modification of $V_0(x)$. The Figure 1 shows one of these first order regular deformations for the simplest Lamé potential

$$V_0(x) = 2m\,\mathrm{sn}^2(x|m). \tag{5}$$

Let us remark that the functions $u(x,-0.1)$, producing the potential $V_1(x,-0.1)$ in Figure 1, preserve the smoothness of (5) as well as its asymptotic behaviour. Moreover, the deformation of V_1 in the neighbourhood of $x=0$ indicates the presence of the bound state energy level at $\epsilon=-0.1$ in the first forbidden gap $(-\infty,E_0)$. For the present case, the related wave-function $1/u(x,-0.1)$ is a smooth function of x, sharply peaked at $x=0$.



FIG. 1: The Lamé potential (5) with $m = 0.5$ (gray) and its first order regular deformation (black) with a bound state energy level at $\epsilon = -0.1 < E_0 = 0.5$ (dashed).

The procedure can be now iterated by considering $\epsilon = \epsilon_a$ as given and $V_1(x,\epsilon_a)$ as the known exactly solvable potential. Thus, one has to look for a new potential $V_2(x,\epsilon_a,\epsilon)$ satisfying the second order relationship

$$V_2(x,\epsilon_a,\epsilon) = V_1(x,\epsilon_a) + 2\beta'(x,\epsilon_a,\epsilon). \tag{6}$$

Therefore, the new function $\beta(x,\epsilon_a,\epsilon)$ must fulfill the Riccati equation $-\beta'(x,\epsilon_a,\epsilon)+\beta^2(x,\epsilon_a,\epsilon) = V_1(x,\epsilon_a)-\epsilon$, where ϵ is arbitrary. The solutions of this equation are easily determined by knowing those of (4) for at least two different factorization constants ϵ, ϵ_a [4]. Then, one can use the finite-difference Bäcklund algorithm [8]. It is remarkable that $\beta(x,\epsilon_a,\epsilon)$ appears to be symmetric under the interchange of ϵ_a by ϵ and vice versa. This means that one can achieve the potential (6) by finding first $V_1(x,\epsilon_a)$ and then $V_2(x,\epsilon_a,\epsilon)$ or, first $V_1(x,\epsilon)$ and then $V_2(x,\epsilon,\epsilon_a)$ [4]. More details concerning further iterations and symmetry properties of the Darboux transformations can be found in [9] and references quoted therein. Observe that potential (6) and the resulting eigenfunctions $\phi_{\epsilon'}(x,\epsilon_a,\epsilon)$ can be rewritten as:

$$V_2(x,\epsilon_a,\epsilon) = V_0(x) - 2\frac{d^2}{dx^2}\ln W(u_a,u);$$

$$\phi_{\epsilon'}(x,\epsilon_a,\epsilon) \propto \frac{W(u_a,u,\psi_{\epsilon'})}{W(u_a,u)}$$

where, for the sake of simplicity, we have written $u_a \equiv u(x,\epsilon_a)$. Let us consider the precedent 1-susy step as given with $\epsilon_a < E_0 \in \sigma(H_0)$. The Hamiltonian $H_1(\epsilon_a)$ is therefore regular and $\phi_{\epsilon'}(x,\epsilon_a,\epsilon)$ becomes an eigenfunction of $H_2(\epsilon_a,\epsilon)$ with eigenvalue ϵ'. If $\epsilon' = E \in \sigma(H_0)$, $\epsilon \neq \epsilon_a$ and $E_0 > \epsilon \in \rho(H_0)$, one easily finds that $\phi_E(x,\epsilon_a,\epsilon)$ can be bounded. Yet, the function $\phi_\epsilon(x,\epsilon_a,\epsilon) \propto 1/\phi_\epsilon(x,\epsilon_a)$ is also orthogonal to all the $\phi_E(x,\epsilon_a,\epsilon)$ and can be chosen in $L^2(\mathbb{R})$. Hence $\sigma(H_2(\epsilon_a,\epsilon)) = \sigma(H_1(\epsilon_a)) \cup \{\epsilon\}$ and the Hamiltonians

$H_1(\epsilon_a)$, $H_2(\epsilon_a, \epsilon)$ are almost isospectral. The iteration chain now establishes that $H_2(\epsilon_a, \epsilon)$ is also a susy partner of H_0 of the second order, *i.e.*, $H_2(\epsilon_a, \epsilon)$ and H_0 are now "sususy partners". In the present case, the iteration procedure leads to nonsingular Hamiltonians at each step, hence, the sususy transformation is said to be *reducible*.



Fig. 2. The initial Lamé potential (gray) and its second order regular deformation (black) allowing two bound state energy levels at $\epsilon_a = 1.1$ and $\epsilon = 1.45$. Observe the asymptotic translational invariance of the potentials.

Now, let us consider the case when $E_0 < \epsilon_a \in \rho(H_0)$. As the function $u(x, \epsilon_a)$ in (5) has a finite number of zeros in $(0, \tau)$, the potential $V_1(x, \epsilon_a)$, as well as the functions $\phi_{\epsilon'}(x, \epsilon_a)$, become singular. However, there is still a chance to construct regular potentials $V_2(x, \epsilon_a, \epsilon)$ and functions $\phi_{\epsilon'}(x, \epsilon_a, \epsilon)$. It is enough to take $\epsilon_a, \epsilon \in \rho(H_0)$ such that $E_j < \epsilon_a, \epsilon < E_{j'}$, with $E_j, E_{j'}$ two subsequent band edges of H_0 (we have used the notation of [7]). Hence, for an appropriate choice of the involved parameters, the Hamiltonian $H_2(\epsilon_a, \epsilon)$ is regular although the intermediary ones $H_1(\epsilon_a)$ and $H_1(\epsilon)$ are singular. We then say that the second order transformation is *irreducible*. Figure 2 shows the result of an irreducible second order regular transformation of potential (5) for $m = 0.5$. The values of the ϵ's are in the gap $(E_1, E_1') = (1, 1.5)$. On the other hand, Figure 3 illustrates one of the related bound states for $\epsilon = 1.45$.



Fig. 3. The probability density for one of the bound states associated to the potential of Figure 2 at $\epsilon = 1.45$

Finally, let us remind that the presence of bound states in the continuum is not a novelty in Quantum Mechanics. For example, Stillinger and Herrick used a modification of the von Neumann-Wigner approach to produce local potentials with bound eigenstates embedded in the continuum of scattering states [10]. Similar results are obtained by means of the susy Quantum Mechanics [11]. However, the situation when the bound states are embedded in the energy gaps of a periodic potential has been out of the attention for researches in the area. Further details on this subject can be found in [5].

The support of CONACyT project 32086E and of CINVESTAV project JIRA'2001/17 is acknowledged.

---

[1] F. Cooper, A. Khare and U. Sukhatme, "Supersymmetry in Quantum Mechanics", World Scientific, Singapore (2000).

[2] B. Mielnik, J. Math. Phys. **25** (1984) 3387; D.J. Fernández C., Lett. Math. Phys. **8** (1984) 337.

[3] C.V. Sukumar, J. Phys. A **19** (1986) 2297; A.A. Andrianov, M.V. Ioffe, F. Cannata and J.P. Dedonder, Int. J. Mod. Phys. A **10** (1995) 6989; V. M. Eleonsky, V.G. Korolev, Phys. Rev. A **55** (19997) 2580; V.G. Bagrov, B.F. Samsonov, Phys. Part. Nucl. **28** (1997) 374; B.F. Samsonov, Phys. Lett. A **263** (1999) 274.

[4] D.J. Fernández C., Int. J. Mod. Phys. A **12** (1997) 171; D.J. Fernández, M.L. Glasser and L.M. Nieto, Phys. Lett. A **240** (1998) 15; J.O. Rosas-Ortiz, J. Phys. A **31** (1998) L507; J. Phys. A **31** (1998) 10163.

[5] D.J. Fernández, B. Mielnik, O. Rosas-Ortiz, B.F. Samsonov, Phys. Lett. A **294** (2002) 168; D.J. Fernández, B. Mielnik, O. Rosas-Ortiz, B.F. Samsonov, J. Phys. A **35** (2002) 4279.

[6] G. Dunne, J. Feinberg, Phys. Rev. D **57** (1998) 1271.

[7] D.J. Fernández, J. Negro, L.M. Nieto, Phys. Lett. A **275** (2000) 338; J. Negro, L.M. Nieto and D.J. Fernández C., Czech. J. Phys. **50** (2000) 1303.

[8] D.J. Fernández C.,V. Hussin and B. Mielnik, Phys. Lett. A **244** (1998) 309; B. Mielnik, L.M. Nieto and O. Rosas-Ortiz, Phys. Lett. A **269** (2000) 70.

[9] L.M. Nieto, B. Mielnik and O. Rosas-Ortiz, in Proceedings of the "VII Encuentro de otoño de Geometría y Física", p. 237, M.A. del Olmo and M. Santander (Eds), Real Sociedad Matemática Española, Madrid, Spain (2001).

[10] F.H. Stillinger and D.R. Herrick, Phys. Rev. A **11** (1975) 446.

[11] J. Pappademos, U. Sukhatme and A. Pagnamenta, Phys. Rev. A **48** (1993) 3525.